

On K_0 -Groups for Substitution Tilings

Johannes Kellendonk

Department of Mathematics, King's College London,
Strand, London WC2R 2LS

E-mail: johannes@math.kcl.ac.uk

Abstract

The group $C(\Omega, \mathbb{Z})/E_{\mathcal{T}}$ is determined for tilings which are invariant under a locally invertible primitive substitution which forces its border. In case the tiling may be obtained by the generalized dual method from a regular grid this group furnishes part of the K_0 -group of the algebra of the tiling. Applied to Penrose tilings one obtains $K_0(\mathcal{A}_{\mathcal{T}}) = \mathbb{Z}^8 \oplus \mathbb{Z}$.

KCL-TH-95-1

Introduction

Non commutative geometry furnishes a framework for the discussion of tilings. Part of the topological aspect of non commutative geometry is the study of the K_0 -group of the algebra of a tiling [1]. This group plays a role for the gap labelling of Schrödinger operators which describe discretized particle systems of solids such as quasicrystals [2, 3] as well as for topological dynamical systems [4, 5].

A tiling is a complete covering of $\mathbb{R}^d$ with distinguished origin 0 by tiles which overlap at most at their boundaries. A tile is closed subset of $\mathbb{R}^d$ carrying a puncture and occasionally more decoration. The punctures of the tiles yield a way to represent a discrete set of allowed translations, i.e. a subset $\mathcal{T}^{pct}$ of $\mathbb{R}^d$. Several tiles form a pattern and we shall be dealing with oriented pattern classes which are equivalence classes of patterns under translations in $\mathbb{R}^d$. Any tiling $\mathcal{T}$ having for any r only finitely many different oriented pattern classes that may be covered by an r -ball defines a compact zero dimensional metric space Ω , its hull, and a (reduced) groupoid- C^* -algebra $\mathcal{A}_{\mathcal{T}}$. The hull is the set of all tilings which are locally homomorphic to $\mathcal{T}$ and such that one of their punctures lies on 0. The groupoid $\mathcal{R} \subset \Omega \times \Omega$ is the groupoid of the equivalence relation $T \sim T'$ whenever T' is a translate of T . We refer for the details, in particular the topological ones, to [6] using the notation of that article partly without further explanation. It is $K_0(\mathcal{A}_{\mathcal{T}})$ together with its natural order structure and scale which is of interest for the gap labelling and as well for d -dimensional topological dynamical systems (defined by d commuting homeomorphisms on a Cantor space) which are related to decorations of $\mathbb{Z}^d$. In the latter case $\mathcal{R}$ is isomorphic as a topological groupoid to a transformation group $\Omega \times_{\varphi} \mathbb{Z}^d$, φ denoting the action of $\mathbb{Z}^d$ on Ω by homeomorphisms, and the algebra is the crossed product $C(\Omega) \times_{\varphi} \mathbb{Z}^d$. The Pimsner Voiculescu six term sequence is the appropriate tool to compute K -groups of such crossed products, and this has been carried out up to $d = 3$ [7]. Using the concept of a reduction of a tiling, which is roughly speaking the same tiling but with a fewer punctures, it could be generalized to tilings (minimal and of dimension smaller or equal 3) which are obtained by the so-called generalized dual method [8] from regular grids [6]. In this case a subset $\mathcal{T}_r^{pct}$ of $\mathcal{T}^{pct}$ could be singled out by taking those punctures which belong to tiles of a chosen oriented pattern class and a continuous action φ of $\mathbb{Z}^d$ on $\Omega_r = \{\mathcal{T} - x | x \in \mathcal{T}_r^{pct}\}$ could be defined so that $C(\Omega_r) \times_{\varphi} \mathbb{Z}^d$ is stably isomorphic to $\mathcal{A}_{\mathcal{T}}$. This includes tilings obtained from the cut and projection method [9] hence in particular the Penrose tilings. To state part of the result recall that for an oriented pattern class M with a puncture of one of its tiles $x \in M^{pct}$

$$\chi_{M,x}(T) = \begin{cases} 1 & \text{if } (M, x) \text{ is at } (T, 0) \\ 0 & \text{else} \end{cases} \quad (1)$$

is a continuous function in the topology of Ω where (M, x) is at $(T, 0)$ means that T contains a pattern of class M such that puncture x coincides with 0. In particular it is a characteristic function on an open and closed set. Moreover the family of those sets generate the topology of the hull so that the above functions generate the group $C(\Omega, \mathbb{Z})$ of continuous functions with integer values. Let $E_{\mathcal{T}}$ be the subgroup which is generated by elements of the form $\chi_{M,x} - \chi_{M,x'}$, $x, x' \in M^{pct}$.

Theorem 1 *Let $\mathcal{T}$ be a minimal tiling obtained by the generalized dual method from a d -dimensional regular grid with $d \leq 3$. Then*

$$\begin{aligned} K_0(\mathcal{A}_{\mathcal{T}}) &= C(\Omega, \mathbb{Z})/E_{\mathcal{T}} && \text{for } d = 1 \\ K_0(\mathcal{A}_{\mathcal{T}}) &= C(\Omega, \mathbb{Z})/E_{\mathcal{T}} \oplus \mathbb{Z} && \text{for } d = 2 \\ K_0(\mathcal{A}_{\mathcal{T}}) &= C(\Omega, \mathbb{Z})/E_{\mathcal{T}} \oplus H && \text{for } d = 3. \end{aligned} \tag{2}$$

Proof: The theorem partly summarizes next to the well known result for $d = 1$ the content of [6]. It is a consequence of $C(\Omega_r) \times_{\varphi} \mathbb{Z}^d$ being stably isomorphic to $\mathcal{A}_{\mathcal{T}}$ and makes use of the results of [7]. The only equality which has still to be shown is $C(\Omega, \mathbb{Z})/E_{\mathcal{T}} = C(\Omega_r, \mathbb{Z})/E_{\varphi}$ where E_{φ} is generated by elements of the form $f - f \circ \varphi(m)$ with $m \in \mathbb{Z}^d$ and $f \in C(\Omega_r, \mathbb{Z})$. It is clear that the inclusion $C(\Omega_r, \mathbb{Z}) \subset C(\Omega, \mathbb{Z})$ induces an embedding of the quotients so that only surjectivity has to be shown. From the minimality of the tiling (which implies that the grid has to be two-sided) follows that there is an R such that whenever an oriented pattern class is large enough in that it covers an R -ball it has to contain a tile which belongs to the chosen oriented pattern class of tiles that carry a puncture in the reduction of the tiling. For M large enough let $x_r \in M^{pct}$ be such a puncture, then $\chi_{M,x} \sim_{E_{\mathcal{T}}} \chi_{M,x_r}$ and $\chi_{M,x_r} \in C(\Omega_r, \mathbb{Z})$. Since $C(\Omega, \mathbb{Z})$ is already generated by functions $\chi_{M,x}$ with large enough patterns the result follows. $\square$

The extra summand $\mathbb{Z}$ for $d = 2$ is related to nontrivial vector bundles over the 2-torus. In fact, $\mathcal{A}_{\mathcal{T}}$ contains the algebra of continuous complex functions over the 2-torus and the generator of the above $\mathbb{Z}$ can be identified with the projection onto a line bundle over it with chern number 1. We do not give the expression for H which is more complicated but known. It explicitly involves the reduction of the tiling.

We will concentrate here on the computation of $C(\Omega, \mathbb{Z})/E_{\mathcal{T}}$ for tilings which are invariant under a locally invertible primitive substitution which forces its border.

1 The group $C(\Omega, \mathbb{Z})/E_{\mathcal{T}}$ for substitution tilings

A substitution of a tiling may be thought of as a rule according to which the tiles of the tiling are to be replaced by patterns which fit together to yield a new tiling. It should, up to rescaling, act covariantly with respect to translation, see [10] for the details. It may as well be understood as a deflation (or generalized decomposition [11]) followed by a rescaling of the tiles to original size. A substitution tiling is a tiling which is invariant up to a translation under a substitution. This expresses a (rather strong) kind of self-similarity. The pattern which replaces a tile or a pattern is called a substitute.

A substitution ρ is locally invertible if one can recognize substitutes by inspection of finite patches (recognizability). This implies that the functions defined for arbitrary oriented pattern classes M and $x \in \rho^n(M)^{pct}$

$$\chi_{M,x}^{(n)}(T) = \begin{cases} 1 & \text{if } (\rho^n(M), x) \text{ is an } n\text{-fold substitute at } (T, 0) \\ 0 & \text{else} \end{cases} \tag{3}$$

are continuous. That $(\rho^n(M), x)$ is an n -fold substitute at $(T, 0)$ means not only that $(\rho^n(M), x)$ is at $(T, 0)$ but that it occurs there as an actual n -fold substitute of M . A locally invertible substitution can be used to construct a continuous map from the hull Ω onto a path space $\mathcal{P}_\Sigma$ over the graph Σ which has the substitution matrix σ as its connectivity matrix. This map is surjective if the substitution is primitive, i.e. if $\rho^n(a)$ will for any tile a and some n contain all other tiles. Furthermore one obtains an embedding $i : \mathcal{A}_\Sigma \rightarrow \mathcal{A}_\mathcal{T}$ of the AF -algebra $\mathcal{A}_\Sigma$ naturally assigned to that path space into the algebra of the tiling. This embedding induces a homomorphism i_* of K -groups which paves the way to compute $K_0(\mathcal{A}_\mathcal{T})$ by determining $K_0(\mathcal{A}_\Sigma)$ as well as $\ker i_*$ and $\operatorname{coker} i_*$. $K_0(\mathcal{A}_\Sigma)$ is computed using the functorial properties of K_0 using the fact that $\mathcal{A}_\Sigma$ is the closure of direct limit of finite dimensional C^* -algebras, see for instance [12]. A substitution forces its border if there is an N such that whenever $\rho^N(a)$ appears in $\mathcal{T}$ as a N -fold substitute its border, i.e. the pattern of tiles which touch it but do not belong to it, are always the same. For simplicity we may assume that $N = 1$ as otherwise we could just work with ρ^N as substitution. The main reason for the introduction of the border forcing condition in [10] was that it guaranteed the mapping from the hull Ω onto the path space $\mathcal{P}_\Sigma$ to be homeomorphic, provided the substitution is primitive. It follows that the tiling is minimal and that $\operatorname{im} i_* = C(\Omega, \mathbb{Z})/E_\mathcal{T}$. The generators of E_Σ are then given by $\chi_{M,x}^{(n)} - \chi_{M,x'}^{(n)}$. In this work we want to compute $\ker i_*$ which is $E_\mathcal{T}/E_\Sigma$, the kernel of the natural projection $C(\Omega, \mathbb{Z})/E_\Sigma \rightarrow (C(\Omega, \mathbb{Z})/E_\Sigma)/(E_\mathcal{T}/E_\Sigma)$.

For an oriented pattern class M we denote by $F(M)$ the oriented pattern class of $\rho(M)$ together with its border, and by $F^n(M)$ the oriented pattern class of $\rho^{n-1}(F(M))$. A 2-pattern is a pattern which consists of two tiles which have a common hypersurface. We denote 2 patterns together with one chosen puncture by $A = (|A|, x)$, i.e. $|A|$ is the corresponding oriented pattern class and $x \in |A|^{pct}$, and write $t(A)$ for the tile which carries the puncture x . $[\cdot]$ indicates equivalence classes with respect to E_Σ .

Lemma 1 *Let $\mathcal{T}$ be invariant under a (primitive) locally invertible substitution which forces its border. Then $E_\mathcal{T}/E_\Sigma$ is generated by elements of the form $[\chi_{M,x}^{(n)}] - [\chi_{M,x'}^{(n)}]$, $x, x' \in \rho^n(M)^{pct}$ where M is a 2 pattern.*

Proof: It is clear that a generating set for $E_\mathcal{T}$ is provided by the set of elements of the form $\chi_{M,x} - \chi_{M,x'}$ where M is arbitrary and $x, x' \in M^{pct}$ are punctures of tiles which have a common hypersurface. Given such M, x, x' consider the set

$$I_{M,x,x'}^{(n)} := \{(A, y) | (M, x) \subset (F^n(|A|), y), y \in \rho^n(t(A))^{pct}, y - x + x' \in \rho^n(|A|)^{pct}\} \quad (4)$$

where $A = (A, x'')$ is a 2 pattern with chosen puncture, $(M, x) \subset (F^n(|A|), y)$ means that the oriented pattern class occurs inside $F^n()$ such that puncture x coincides with y , and we keep track of the position of $\rho^n(t(A))$ and $\rho^n(|A|)$ in $F^n(|A|)$ to whether a puncture of $F^n(|A|)$ lies in $\rho^n(t(A))$ or $\rho^n(|A|)$. Since ρ determines its border we have for big enough n

$$[\chi_{M,x}] - [\chi_{M,x'}] = \sum_{(A,y) \in I_{M,x,x'}^{(n)}} [\chi_{|A|,y}^{(n)}] - [\chi_{|A|,y-x+x'}^{(n)}] \quad (5)$$

which proves the lemma. $\square$

Using the shorter notation $\chi_i^{(n)} := [\chi_{a_i, x}^{(n)}]$ with some $x \in \rho^n(a_i)^{pct}$ and $\chi_A^{(n)} := [\chi_{|A|, x}^{(n)}]$ with $x \in \rho^n(t(A))^{pct}$ the group $C(\Omega, \mathbb{Z})/E_\Sigma$ is generated by $\{\chi_i^{(n)}\}_i$ and the result of the above lemma can be formulated by saying that $E_\mathcal{T}/E_\Sigma$ is generated by elements of the form $\chi_A^{(n)} - \chi_{\bar{A}}^{(n)}$ where $\bar{A} = (A, \bar{x})$ and $\bar{x}$ is the puncture of the tile which is not $t(A)$. Recall the definition of the substitution matrix

$$\sigma_{ij} = \text{number of } a_j \text{'s in } \rho(a_i) \quad (6)$$

which satisfies

$$\chi_j^{(n)} = \sum_i \chi_i^{(n+1)} \sigma_{ij}. \quad (7)$$

In a similar spirit let

$$K_{iA} := \text{number of } A \text{'s in } F(a_i) \text{ with } t(A) \text{ in } \rho(a_i) \quad (8)$$

for which we keep track of the position of $\rho(a)$ in $F(a)$. Then

$$\chi_A^{(n)} = \sum_i \chi_i^{(n+1)} K_{iA}. \quad (9)$$

Define

$$\mathcal{K}_{iA} := K_{iA} - K_{i\bar{A}} \quad (10)$$

and $\mathcal{K}^{(n)} : \mathbb{Z}^s \rightarrow C(\Omega, \mathbb{Z})/E_\Sigma : e_A \mapsto \sum_i \chi_i^{(n)} \mathcal{K}_{iA}$, where s is the number of 2 pattern classes together with a chosen puncture. Note that the matrix $\sigma^n \mathcal{K}$ plays the role of $\mathcal{K}$ for the substitution ρ^{n+1} , $n \geq 0$.

Theorem 2 *Let $\mathcal{T}$ be a tiling with locally invertible substitution which forces its border. Then $E_\mathcal{T}/E_\Sigma$ is generated by $\bigcup_n \text{im } \mathcal{K}^{(n)}$. If moreover the substitution matrix σ is invertible over $\mathbb{Z}$ so that $C(\Omega, \mathbb{Z})/E_\Sigma \cong \mathbb{Z}^r$ then*

$$C(\Omega, \mathbb{Z})/E_\mathcal{T} = \mathbb{Z}^r / < \bigcup_{n \leq 0} \text{im } \sigma^n \mathcal{K} > \quad (11)$$

where $\mathcal{K} : \mathbb{Z}^s \rightarrow \mathbb{Z}^r : \mathcal{K}(e_A)_i = \mathcal{K}_{iA}$.

Proof: Using (7) and the last lemma the classes of the generators of $E_\mathcal{T}/E_\Sigma$ may be expressed as

$$\chi_A^{(n)} - \chi_{\bar{A}}^{(n)} = \sum_i \chi_i^{(m)} (\sigma^{m-n-1} \mathcal{K})_{iA} \quad (12)$$

where m is any integer which has, in case that σ is not invertible to be greater or equal to $n+1$. Taking $m = n+1$ one obtains $E_\mathcal{T}/E_\Sigma = < \bigcup_n \text{im } \mathcal{K}^{(n)} >$. If σ is invertible we may take $m = 0$ which implies that $E_\mathcal{T}/E_\Sigma$ is generated by $\bigcup_{n \leq 0} \text{im } \sigma^n \mathcal{K}$. $\square$

Note that due to the subtraction in the definition of $\mathcal{K}$ above one only has to count those 2 patterns in $F(a)$ which cross the boundary of $\rho(a)$ and take into account only one choice for the puncture in $|A|$.

2 Applications

Before we apply Theorem 2 to Penrose tilings let us have a look at the one dimensional situation (substitution sequences) and show that (11) fits into the more general results of [13]. Taking a suitable power of the substitution, which we suppose to be locally invertible primitive and to force its border, we may as well assume that the first and last letters of substitutes are stable under it, i.e. that $\rho(\rho(a)_f)_f = \rho(a)_f$ resp. $\rho(\rho(a)_l)_l = \rho(a)_l$, where w_f, w_l denote the first resp. last letter of the word w . Let us denote by $\mathcal{F}$ the indices of the first letters and by $\mathcal{L}$ those of the last. The border forcing condition implies that there is a bijection $\varphi : \mathcal{L} \rightarrow \mathcal{F}$. This means that whenever a_j is the last (resp. first) letter of $\rho(a_i)$ then the letter following (resp. standing ahead of) it in the sequence is $a_{\varphi(j)}$ (resp. $a_{\varphi^{-1}(j)}$). To evaluate (11) we may restrict the consideration to the 2 patterns $a_j a_{\varphi(j)}$, $j \in \mathcal{L}$ and choose the puncture to belong to a_j ; we abbreviate them as $[j\varphi(j)]$. This leads to consider $\mathcal{K} : \mathbb{Z}^{|\mathcal{L}|} \rightarrow \mathbb{Z}^r$, r being the total number of letters, which has entries

$$\mathcal{K}_{ij} := \mathcal{K}_{i[j\varphi(j)]} = \begin{cases} 0 & \text{if } a_j \neq \rho(a_i)_l \text{ and } a_{\varphi(j)} \neq \rho(a_i)_f \\ 0 & \text{if } a_j = \rho(a_i)_l \text{ and } a_{\varphi(j)} = \rho(a_i)_f \\ 1 & \text{if } a_j = \rho(a_i)_l \text{ and } a_{\varphi(j)} \neq \rho(a_i)_f \\ -1 & \text{if } a_j \neq \rho(a_i)_l \text{ and } a_{\varphi(j)} = \rho(a_i)_f \end{cases} \quad (13)$$

$j \in \mathcal{L}$. Since $\sigma\mathcal{K}$ plays the role of $\mathcal{K}$ but for ρ^2 the above stability of first and last letters under substitution implies

$$\mathcal{K} = \sigma\mathcal{K}. \quad (14)$$

Now let us take a piece of the tiling $\rho(a_{i_1})\rho(a_{i_2})\cdots\rho(a_{i_n})$ and obtain from it a sequence of last letters $j_1 j_2 \cdots j_n$, $j_k = \rho(a_{i_k})_l$. If n is large enough this sequence will by primitivity of the substitution contain all $j \in \mathcal{L}$. But two consecutive elements $j_{k-1} j_k$ of the sequence indicate that (in vector notation) $\mathcal{K}_{i_k} = e_{j_k} - e_{j_{k-1}}$ so that the occurrence of all $j \in \mathcal{L}$ implies that $\text{im } \mathcal{K}$ is generated by $\{e_j - e_{j'}\}_{j, j' \in \mathcal{L}}$. Hence in case σ is invertible we get $\text{coker } \mathcal{K} = \mathbb{Z}^{r-|\mathcal{L}|+1}$.

2.1 $K_0(\mathcal{A}_{\mathcal{T}})$ for Penrose tilings

There are several well known variants of tilings which are called Penrose tilings and which are a priori to be distinguished as they lead to non-isomorphic groupoids. The version which is most suitable for our purposes is the one which has triangles as tiles, cf. Figure 2. The triangles are decorated (with a little circle) to break the mirror symmetry. We may take the centers of these circles to be the punctures. These Penrose tilings by triangles possess an orientational symmetry of 20 elements which is generated by a rotation around $\frac{\pi}{5}$ together with a mirror reflection at a boundary line of a triangle [10]. A Penrose tiling by triangles has 40 oriented pattern classes of tiles. A short glance at such a tiling shows that always two triangles of the same congruence class (under all Euclidian transformations) yield a 2 pattern which is mirror symmetric and forms a rhombus. Deleting the corresponding lines yields a tiling made from rhombi. As a consequence, such a Penrose tiling by rhombi has only 20 oriented pattern classes of

tiles. A Penrose tiling by rhombi is actually a reduction of a Penrose tiling by triangles. Let $\mathcal{T}_{Rh}^{pct}$ consist of all punctures which belong to tiles that are obtained from those of Figure 2.1 and 2.3 by a rotation. This clearly defines a reduction and consequently the algebra of a Penrose tiling by rhombi is stably isomorphic to the one of a Penrose tiling by triangles and the corresponding K_0 -groups are isomorphic as ordered groups. In fact, like in the proof of Theorem 1 one verifies that $C(\Omega_{Rh}, \mathbb{Z})/E_{\mathcal{T}_{Rh}} = C(\Omega, \mathbb{Z})/E_{\mathcal{T}}$ where $E_{\mathcal{T}_{Rh}} = E_{\mathcal{T}} \cap C(\Omega_{Rh}, \mathbb{Z})$. We prefer to work with Penrose tilings by triangles since they allow for a substitution which is covariant under the orientational symmetry, and this may be used to simplify the computation of $\text{im } \mathcal{K}$.

Those tilings which have an exact five-fold symmetry are invariant under the substitution $\rho = \check{\rho}^4$ the deflation corresponding to $\check{\rho}$ being displayed in Figure 1 for one orientation.




Figure 1

This substitution ρ is primitive, locally invertible, and forces its border with $N = 1$ [10]. Its substitution matrix is invertible so that $C(\Omega, \mathbb{Z})/E_{\Sigma} = \mathbb{Z}^{40}$. To simplify the computation of $E_{\mathcal{T}}/E_{\Sigma}$ we make use of the symmetry properties of the tiling and in particular of the fact that 2 patterns crossing the boundaries of $\rho(a)$ are always mirror symmetric. Let α denote the direction of a boundary line of a tile resp. a substitute. There are 10 different ones and we order them anti-clockwise identifying them with $\{0, \dots, 9\}$. Saying that a substitute has boundary α if it has a boundary with that direction we define

$$N_{ij}^{\alpha} := \text{number of } a_j \text{'s at boundary } \alpha \text{ of } \rho(a_i). \quad (15)$$

In particular $N_{ij}^{\alpha} = 0$ in case a_j or $\rho(a_i)$ do not have boundary α . Let $a_{\alpha(j)}$ be the mirror image of a_j with respect to the mirror axis α and define

$$\mathcal{N}_{ij}^{\alpha} = N_{ij}^{\alpha} - N_{i\alpha(j)}^{\alpha} \quad (16)$$

Then setting $\mathcal{N} : \mathbb{Z}^{10} \otimes \mathbb{Z}^{40} \rightarrow \mathbb{Z}^{40} : \mathcal{N}(e_{\alpha} \otimes e_j)_i = \mathcal{N}_{ij}^{\alpha}$ we have

$$\mathcal{K}_{iA} = \begin{cases} \mathcal{N}_{ij}^{\alpha} & \text{if } |A| = a_j a_{\alpha(j)} \text{ and } t(A) = a_j \\ -\mathcal{N}_{ij}^{\alpha} & \text{if } |A| = a_j a_{\alpha(j)} \text{ and } t(A) = a_{\alpha(j)} \\ 0 & \text{else} \end{cases} \quad (17)$$

$|A| = a_j a_{\alpha(j)}$ indicating that $|A|$ is composed of a_j and $a_{\alpha(j)}$. Hence $\text{im } \mathcal{K} = \text{im } \mathcal{N}$. $\mathcal{N}^\alpha$ is related to $\mathcal{N}^0$ by symmetry, i.e. $\mathcal{N}^\alpha = R^{-\alpha} \mathcal{N}^0 R^\alpha$, R being the matrix which acts as a rotation around $\frac{\pi}{5}$. Let us use a basis $\{\chi_{10k+\alpha}^{(0)}\}_{0 \leq k \leq 3, 0 \leq \alpha \leq 9}$ in which $a_{10k+\alpha}$ corresponds to the oriented pattern class of the triangle in Figure 2.k rotated around an angle of $\frac{\alpha\pi}{5}$.




Figure 2.0




Figure 2.1



Figure 2.2



Figure 2.3

In terms of the rotation matrix ω having entries $\omega_{\alpha\beta} = \delta_{\beta-\alpha, 1 \bmod 10}$ R and the substitution matrix $\sigma = \check{\sigma}^4$ are given by

$$R = \begin{pmatrix} \omega & 0 & 0 & 0 \\ 0 & \omega & 0 & 0 \\ 0 & 0 & \omega & 0 \\ 0 & 0 & 0 & \omega \end{pmatrix} \quad (18)$$

$$\check{\sigma} = \begin{pmatrix} \omega^4 & \omega^0 & 0 & \omega^6 \\ \omega^0 & \omega^6 & \omega^4 & 0 \\ \omega^3 & 0 & \omega^7 & 0 \\ 0 & \omega^7 & 0 & \omega^3 \end{pmatrix}. \quad (19)$$

The matrix N^0 may be read of from Figure 3 (and completed by symmetry); it is given below (27). Moreover $\mathcal{N}^0 = N^0 - N^0 S$ where S implements the reflection at $\alpha = 0$, explicitly, with $s_{\alpha\beta} = \delta_{\alpha+\beta, 5 \bmod 10}$ (counting rows and columns form 0 to 9)

$$S = \begin{pmatrix} 0 & s & 0 & 0 \\ s & 0 & 0 & 0 \\ 0 & 0 & 0 & s \\ 0 & 0 & s & 0 \end{pmatrix}. \quad (20)$$

It turns out that $\text{im } \mathcal{N}$ is generated by the orbit under R of the four vectors

$$\begin{aligned} v_1 &= (0,1,0,0,0,0,-1,0,0,0,0,0,0,0,0) \\ v_2 &= (0,0,0,0,0,0,1,0,0,0,0,-1,0) \\ w_1 &= (0,0,0,0,0,0,0,1,0,0,-1,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,1,0,0,0,0,0,0,0,-1) \\ w_2 &= (0,0,0,0,0,1,0,0,0,0,0,-1,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,-1,0,0,0,0,0,1,0) \end{aligned} \quad (21)$$

but 8 of the 40 vectors thus obtained are linearly dependent. Moreover $\text{im } \mathcal{N}$ is invariant under σ . Dividing it out yields no torsion so that we obtain

$$K_0(\mathcal{A}_{\mathcal{T}}) \cong \mathbb{Z}^8 \oplus \mathbb{Z}. \quad (22)$$

Let us interpret this result. Let H_1 be the sublattice of $\mathbb{Z}^{40}$ which is generated by $\{R^\alpha v_i\}_{i=1,2; \alpha=0,1,\dots,9}$. $(C(\mathcal{P}_\Sigma, \mathbb{Z})/E_\Sigma)/H_1 = \mathbb{Z}^{20}$ has a basis with natural geometrical interpretation. It is formed by the characteristic functions on the oriented pattern classes of rhombi which are obtained from the triangles as described above.

The Penrose tilings by rhombi as they are given by reducing Penrose tilings by triangles as above do not have symmetric tiles. We may delete the circle in the center of which there is no puncture. This reduces also the orientational symmetry to only 10 elements (the mirror reflection interchanges punctures from $\mathcal{T}_{Rh}^{pct}$ with those from $\mathcal{T}^{pct} \setminus \mathcal{T}_{Rh}^{pct}$). But to our knowledge there is no substitution for Penrose tilings by rhombi which is covariant under the orientational symmetry. But there are at least 10 non covariant ones; namely $\rho_\alpha = \check{\rho}_\alpha^4$ the deflation corresponding to $\check{\rho}_0$ being given in Figure 4 and $\check{\rho}_\alpha$ being obtained from $\check{\rho}_0$ just by rotation of the whole figure around $\frac{\alpha\pi}{5}$, $\alpha = 0, \dots, 9$. That all these substitutions are primitive locally invertible and force their border carries over from ρ . In the basis corresponding to the rhombi ordered as in Figure 4 from above to below the substitution matrix of $\check{\rho}_0$ is given by

$$\check{\sigma}_0 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}. \quad (23)$$

It turns out that

$$\ker \check{\sigma}_\alpha = \langle \{R^\alpha w_i\}_{i=1,2} \rangle. \quad (24)$$

Since the restriction to its image is an automorphism and $\ker \check{\sigma}_0 = \ker \check{\sigma}_0^2$ we have

$$C(\mathcal{P}_{\Sigma_\alpha}, \mathbb{Z})/E_{\Sigma_\alpha} \cong \mathbb{Z}^{20} / \langle \{R^\alpha w_i\}_{i=1,2} \rangle \quad (25)$$

Σ_α denoting the graph having $\tilde{\sigma}_\alpha$ as connectivity matrix. Let $j : \mathbb{Z}^{20} \rightarrow C(\Omega_{Rh}, \mathbb{Z})/E_{\mathcal{T}_{Rh}}$ be given by assigning the characteristic functions onto the oriented pattern classes of rhombi to the standard basis. It is surjective since $H_1 \subset \text{im } \mathcal{N}$. Any of the substitutions ρ_α leads to a homeomorphism between Ω_{Rh} and the path space $\mathcal{P}_{\Sigma_\alpha}$ defined by Σ_α and moreover to an embedding $i_\alpha : \mathcal{A}_{\Sigma_\alpha} \rightarrow \mathcal{A}_{\mathcal{T}}$ of the AF -algebra naturally assigned to $\mathcal{P}_{\Sigma_\alpha}$ into the algebra of $\mathcal{T}$. Let π_α be the natural projection onto $\mathbb{Z}^{20}/\ker \sigma_\alpha$ then $j = (i_\alpha)_* \circ \pi_\alpha$ for all α and therefore

$$\ker j \subset \langle \{R^\alpha w_i\}_{i=1,2; \alpha=1,\dots,9} \rangle \cong \mathbb{Z}^{12}. \quad (26)$$

This shows independently that $C(\Omega, \mathbb{Z})/E_{\mathcal{T}} \subset \mathbb{Z}^8$ but not the opposite inclusion. A computation of $\text{im } \mathcal{K}$ for e.g. ρ_0 would have been more complicated due to the lack of symmetry.

It remains to compute the order structure. The above result for the K_0 -group raises the question whether there is a tiling which is mutually locally derivable from a Penrose tiling but has only 8 oriented pattern classes of tiles.

$$N^0 = \begin{pmatrix} 3000000000100010200000000001001000000000 \\ 0100000001000002000000000100000000000010 \\ 00 \\ 00 \\ 1000101000100000100010000000000000000000 \\ 0000030000020001000100100000000000010000 \\ 0000101000200000000010000000000001000000 \\ 00 \\ 00 \\ 0100010001010001000000000100000000000000 \\ 1000201000300000000010000000000001000000 \\ 0000010001010001000100000000000000010000 \\ 00 \\ 00 \\ 20000000000000010100000000001001000000000 \\ 0100010002000003000000000100000000000010 \\ 1000100000100010100000000000000100000000 \\ 00 \\ 00 \\ 0000020000010000000100100000000000010000 \\ 10000000000000010100000000000000100000000 \\ 00 \\ 0100000001000002000000000100000000000010 \\ 1000100000100010100000000000000100000000 \\ 00 \\ 0000010000010000000100000000000000010000 \\ 00 \\ 0000101000200000000010000000000001000000 \\ 0000010001010001000100000000000000010000 \\ 00 \\ 0000101000100000000010000000000000000000 \\ 00 \\ 0100010001010001000000000100000000000000 \\ 2000000000000001010000000000100100000000 \\ 00 \\ 0100000001000001000000000100000000000000 \\ 00 \\ 1000101000100000100010000000000000000000 \\ 0000020000010000000100100000000000010000 \\ 00 \end{pmatrix} \quad (27)$$




Figure 3

Substitutes of $\rho = \check{\rho}^4$ of the triangles of Figure 2.0 and 2.1 in all directions.




Figure 4

Acknowledgement. I thank Alan Forrest for very fruitful conversations.

References

- [1] A. Connes. *Géométrie Non Commutative*. InterEditions (Paris), 1990.
- [2] J. Bellissard. K -Theory of C^* -Algebras in Solid State Physics. In T.C. Dorlas, N.M. Hugenholtz, and M. Winnik, editors, *Statistical Mechanics and Field Theory: Mathematical Aspects*. Lect. Notes in Phys. 257:99-156, Springer Verlag, 1986.
- [3] J. Bellissard, A. Bovier, and J.M. Ghez. Gap labelling theorems for one dimensional discrete Schrödinger operators. *Rev. Math. Phys.*, 4:1–38, 1992.
- [4] R.H. Herman, I.F. Putnam, and C.F. Skau. Ordered Bratteli diagrams, dimension groups and topological dynamics. *Int. J. Math.*, 3:827–864, 1992.
- [5] T. Giordano, I.F. Putnam, and C.F. Skau. Topological orbit equivalence and C^* -crossed products. Trondheim preprint Mathematics No. 11/1993, 1993.
- [6] J. Kellendonk. Gap labelling for Schrödinger operators on quasiperiodic tilings. King’s College London preprint KCL-TH-94-10, cond-mat 9406107, 1994.
- [7] A. van Elst. Gap-labelling theorems for Schrödinger operators on the square and cubic lattice. *Rev. Math. Phys.*, 6:319–342, 1994.
- [8] J.E.S. Socolar, P.J. Steinhardt, and D. Levine. Quasicrystals with arbitrary orientational symmetry. *Phys. Rev. B*, 32:5547–5550, 1985.
- [9] M. Duneau and A. Katz. Quasiperiodic patterns and icosahedral symmetry. *J. Physique*, 47:181–196, 1986.
- [10] J. Kellendonk. Non commutative geometry of tilings and gap labelling. King’s College London preprint KCL-TH-94-1, cond-mat 9403065, 1994.
- [11] B. Grünbaum and G.C. Shephard. *Tilings and Patterns*. Freeman and Company (New York), 1987.
- [12] G.J. Murphy. *C^* -Algebras And Operator Theory*. Academic Press, 1990.
- [13] A.H. Forrest. K -groups associated with substitution minimal systems. Edinburgh preprint 1994, submitted to Israel J. Math.